

Quantum Zeno effect and the impact of flavor in leptogenesis

S. Blanchet, P. Di Bari and G. G. Raffelt

Max-Planck-Institut für Physik (Werner-Heisenberg-Institut)
Föhringer Ring 6, 80805 München, Germany

MPP-2006-155 (17 November 2006)

Abstract. In thermal leptogenesis, the cosmic matter-antimatter asymmetry is produced by CP violation in the decays $N \rightarrow \ell + \Phi$ of heavy right-handed Majorana neutrinos N into ordinary leptons ℓ and Higgs particles Φ . If some charged-lepton Yukawa couplings are in equilibrium during the leptogenesis epoch, the ℓ interactions with the background medium are flavor sensitive and the coherence of their flavor content defined by $N \rightarrow \ell + \Phi$ is destroyed, modifying the efficiency of the inverse decays. We point out, however, that it is not enough that the flavor-sensitive processes are fast on the cosmic expansion time scale, they must be fast relative to the $N \leftrightarrow \ell + \Phi$ reactions lest the flavor amplitudes of ℓ remain frozen by the repeated $N \leftrightarrow \ell + \Phi$ “measurements”. Our more restrictive requirement is significant in the most interesting “strong wash-out case” where $N \leftrightarrow \ell + \Phi$ is fast relative to the cosmic expansion rate. We derive conditions for the unflavored treatment to be adequate and for flavor effects to be maximal. In this “fully flavored regime” a neutrino mass bound survives. To decide if this bound can be circumvented in the intermediate case, a full quantum kinetic treatment is required.

1. Introduction

The see-saw mechanism [1–6] is an elegant way to understand neutrino masses and mixing and at the same time, with leptogenesis [7], provides a natural mechanism for generating the matter-antimatter asymmetry of the Universe by virtue of CP -violating decays $N \rightarrow \ell + \Phi$ of heavy right-handed Majorana neutrinos N into ordinary leptons ℓ and Higgs particles Φ . In comparison with other baryogenesis scenarios, leptogenesis has the unique advantage of relying on an ingredient of Physics Beyond the Standard Model, neutrino masses, that is experimentally established. Moreover, the neutrino mass values suggested by flavor oscillation experiments are optimal in the sense that leptogenesis would operate in a mildly “strong wash-out regime” [8]. This means that the inverse processes $\ell + \Phi \rightarrow N$ are efficient enough to wash out all contributions to the final asymmetry that depend on initial conditions, but not too efficient to prevent successful leptogenesis. This is true for hierarchical light neutrino schemes, while in the case of quasi-degenerate neutrinos, leptogenesis provides a stringent upper bound

on neutrino masses, $m_1 \leq 0.1 \text{ eV}$, holding when a hierarchical right-handed neutrino spectrum is assumed [8].

Traditionally, flavor effects were neglected in leptogenesis, i.e., the lepton asymmetry produced by $N \rightarrow \ell + \Phi$ and the wash-out effects caused by the inverse processes were treated as if ℓ had no flavor properties. Recently it was recognized, however, that important modifications arise if some of the charged-lepton Yukawa interactions are in equilibrium [9–12]. In particular, there is an additional source of CP violation that can make the single-flavored CP asymmetries larger than the total. This additional source includes a dependence on low-energy phases, in contrast to unflavored leptogenesis [10, 11]. In fact, successful leptogenesis stemming only from Majorana and Dirac phases is possible [13–16]. A second consequence is a reduction of the wash-out efficiency, generally by different amounts in each flavor. These modifications together contribute to enhance the final asymmetry, an effect that increases for a larger neutrino mass scale. Therefore, flavor effects can relax the neutrino mass bound of the unflavored scenario [12].

The purpose of our present study is to explore the conditions for flavor effects to be relevant in leptogenesis and if the neutrino mass bound can indeed be circumvented. The past literature is unclear on this point in the following sense. It was stressed that the ℓ interactions with the background medium are flavor sensitive and the coherence of the ℓ flavor content defined by $N \rightarrow \ell + \Phi$ can be destroyed if some of the charged-lepton Yukawa couplings are in thermal equilibrium during the leptogenesis epoch. On the other hand, the reactions $N \leftrightarrow \ell + \Phi$ constantly re-generate the ℓ flavor composition and if these reactions are fast, ℓ is frozen in its coherent flavor state by the quantum Zeno effect [17, 18]. In other words, the charged-lepton Yukawa couplings and the $N \leftrightarrow \ell + \Phi$ reactions generally single out different directions in flavor space. The ℓ density matrix in flavor space will dominantly stay in the direction favored by the fastest term of the kinetic equation. Therefore, we expect flavor effects to be important if the “flavor measurements” by the background medium are fast relative to the $N \leftrightarrow \ell + \Phi$ reactions, a condition that was also stated in Ref. [11]. On the other hand, the actually used requirement in the previous literature was the weaker condition that the flavor-sensitive reactions are fast on the cosmic expansion scale. Of course, at the end of the leptogenesis epoch when the $N \leftrightarrow \ell + \Phi$ reactions no longer track equilibrium, this weaker condition is correct. Our main concern is that previous treatments may not properly cover the important strong wash-out case, where $N \leftrightarrow \ell + \Phi$ is fast relative to the expansion rate for most of the leptogenesis epoch.

In order to identify the conditions for flavor effects to be relevant for the final baryon asymmetry we split the problem into two parts. First, in Section 2, we derive a condition for the unflavored treatment to be adequate. Next, in Section 3, we obtain a condition for flavor effects to be maximally effective (“fully flavored regime”). In these cases two different sets of classical Boltzmann equations apply to calculate the final asymmetry. In Section 4 we consider the neutrino mass bound to hold in the unflavored case and show that it cannot be circumvented in the fully flavored case. However,

there is an intermediate regime between the fully flavored and unflavored cases where a full quantum kinetic equation is required to decide on the neutrino mass bound. In Section 5 we briefly discuss the limitations of our simple rate comparison. We summarize our findings in Section 6.

2. Unflavored case

2.1. Unflavored leptogenesis

In order to derive a condition for the unflavored leptogenesis treatment to be adequate, we first retrace the steps leading to the final asymmetry prediction. Adding to the Standard Model three right-handed (RH) neutrinos with a Majorana mass term M and Yukawa couplings h , after spontaneous symmetry breaking a Dirac mass term, $m_D = v h$, is generated by the vev v of the Higgs boson. In the see-saw limit, $M \gg m_D$, the spectrum of neutrino masses splits into two sets, a very heavy one, $M_3 \geq M_2 \geq M_1$, almost coinciding with the eigenvalues of M , and a light one, $m_3 \geq m_2 \geq m_1$, corresponding to the eigenvalues of the light neutrino mass matrix. It is given by the see-saw formula [1–6],

$$m_\nu = -m_D \frac{1}{M} m_D^T. \quad (1)$$

Neutrino oscillation experiments measure two light neutrino mass squared differences. In a normal scheme one has $m_3^2 - m_2^2 = \Delta m_{\text{atm}}^2$ and $m_3^2 - m_1^2 = \Delta m_{\text{sol}}^2$, whereas in an inverted scheme one has $m_3^2 - m_2^2 = \Delta m_{\text{sol}}^2$ and $m_2^2 - m_1^2 = \Delta m_{\text{atm}}^2$. For $m_1 \gg m_{\text{atm}} \equiv \sqrt{\Delta m_{\text{atm}}^2}$ the spectrum is quasi-degenerate, while for $m_1 \ll m_{\text{sol}} \equiv \sqrt{\Delta m_{\text{sol}}^2}$ it is fully hierarchical.

In the early Universe, the decays of the RH neutrinos into leptons and Higgs bosons generally produce a net lepton number that is partly converted into a net baryon number by sphaleron processes if the temperature exceeds about 100 GeV. We will follow common practice and assume that the final asymmetry is determined only by the decays of the lightest RH neutrino, although a scenario where the asymmetry is produced by the next-to-lightest states is also possible [19, 20].

For now we also assume that the flavor composition of the leptons produced in the decays does not affect the final asymmetry (unflavored limit). Neglecting thermal effects in $\Delta L = 1$ scatterings [21] and spectator processes [22, 23], the final asymmetry is determined by a simple set of two kinetic equations, one for the RH neutrino abundance and one for the $B - L$ asymmetry [24],

$$\frac{dN_{N_1}}{dz} = -D \left(N_{N_1} - N_{N_1}^{\text{eq}} \right) \quad (2)$$

$$\frac{dN_{B-L}}{dz} = \varepsilon_1 D \left(N_{N_1} - N_{N_1}^{\text{eq}} \right) - \left(W_1^{\text{ID}} + \Delta W \right) N_{B-L}, \quad (3)$$

where $z \equiv M_1/T$. With N_{B-L} and N_{N_1} we denote the abundances per RH neutrino N_1 , taken to be in ultra-relativistic thermal equilibrium. The actual N_1 equilibrium abundance is $N_{N_1}^{\text{eq}} = z^2 \mathcal{K}_2(z)/2$, where $\mathcal{K}_i(z)$ are the modified Bessel functions.

Introducing the decay parameter $K_1 \equiv \tilde{\Gamma}_1/H_{T=M_1}$, defined as the ratio of the total decay width to the expansion rate at $T = M_1$, the decay term can be written as

$$D \equiv \frac{\Gamma_D}{H z} = K_1 z \langle \gamma_1^{-1} \rangle, \quad (4)$$

where $\langle \gamma_1^{-1} \rangle = \mathcal{K}_1(z)/\mathcal{K}_2(z)$ is the thermally averaged Lorentz dilation factor. The expansion rate is

$$H \simeq \sqrt{\frac{8\pi^3 g_\star}{90}} \frac{M_1^2}{M_{\text{Pl}}} \frac{1}{z^2} \simeq 1.66 \sqrt{g_\star} \frac{M_1^2}{M_{\text{Pl}}} \frac{1}{z^2} \quad (5)$$

and $g_\star = g_{\text{SM}} = 106.75$ is the total number of thermally excited degrees of freedom.

The wash-out term caused by inverse decays is, after subtracting the resonant $\Delta L = 2$ processes,

$$W_1^{\text{ID}}(z) \equiv \frac{1}{2} \frac{\Gamma_1^{\text{ID}}(z)}{H(z) z} = \frac{1}{4} K_1 \mathcal{K}_1(z) z^3, \quad (6)$$

where $\Gamma_1^{\text{ID}} = \Gamma_D N_{N_1}^{\text{eq}}/N_\ell^{\text{eq}}$ is the inverse decay rate. The non-resonant $\Delta L = 2$ contribution dominates in the non-relativistic regime and is approximately

$$\Delta W(z) \simeq \frac{w}{z^2} \left(\frac{M_1}{10^{10} \text{ GeV}} \right) \left(\frac{\bar{m}}{\text{eV}} \right)^2, \quad (7)$$

where

$$w = \frac{9\sqrt{5} M_{\text{Pl}} 10^{-8} \text{ GeV}^3}{4\pi^{9/2} g_\ell \sqrt{g_\star} v^4} \simeq 0.186 \quad (8)$$

and $\bar{m} \equiv \sqrt{\sum_i m_i^2}$.

The evolution of N_{B-L} is explicitly, denoting initial quantities with the subscript “in,”

$$N_{B-L}(z) = N_{B-L}^{\text{in}} \exp \left\{ - \int_{z_{\text{in}}}^z dz' [W_1^{\text{ID}}(z') + \Delta W(z')] \right\} + \varepsilon_1 \kappa_1(z) \quad (9)$$

with the efficiency factor

$$\kappa_1(z) \equiv - \int_{z_{\text{in}}}^z dz' \frac{dN_{N_1}}{dz'} \exp \left[- \int_{z'}^z dz'' [W_1^{\text{ID}}(z'') + \Delta W(z'')] \right]. \quad (10)$$

An approximate analytic expression for the final efficiency factor at the end of the leptogenesis epoch and valid in the strong wash-out regime ($K_1 \gg 1$) is

$$\kappa_1^{\text{f}}(m_1, M_1, K_1) \simeq \kappa(K_1) \exp \left[- \frac{w}{z_B(K_1)} \left(\frac{M_1}{10^{10} \text{ GeV}} \right) \left(\frac{\bar{m}}{\text{eV}} \right)^2 \right] \quad (11)$$

where,

$$\kappa(K_1) \simeq \frac{2}{K_1 z_B(K_1)} \left(1 - e^{-K_1 z_B(K_1)/2} \right). \quad (12)$$

Further,

$$z_B(K_1) \simeq 2 + 4 K_1^{0.13} e^{-2.5/K_1} \quad (13)$$

is an approximate expression for that value of z where $W_1^{\text{ID}}(z_B) \simeq 1$, i.e., where the wash-out term from inverse decays becomes ineffective [25]. In the weak wash-out regime

the expression for the final efficiency factor depends on the value of $N_{N_1}^{\text{in}}$. For an initial thermal equilibrium abundance the given expression still holds. For an initial vanishing abundance the final efficiency factor is the sum of two contributions with opposite sign and approximate analytic expressions can be found in [24]. The weak wash out, acting less efficiently on the positive contribution, prevents a full cancellation.

Assuming either that the initial asymmetry is negligible or that it is efficiently washed out, the final $B - L$ asymmetry is $N_{B-L}^f \simeq \varepsilon_1 \kappa_1^f$. Assuming further a standard thermal history of the universe and accounting for the sphaleron conversion coefficient $a_{\text{sph}} \sim 1/3$, the final baryon-to-photon ratio at recombination (rec) is

$$\eta_B = a_{\text{sph}} \frac{N_{B-L}^f}{N_{\gamma}^{\text{rec}}} \simeq 0.96 \times 10^{-2} \varepsilon_1 \kappa_1^f. \quad (14)$$

This is the number to be compared with the measured value [26]

$$\eta_B^{\text{CMB}} = (6.1 \pm 0.2) \times 10^{-10}. \quad (15)$$

The wash-out rate by inverse decays reaches a maximum $W_1^{\text{ID}}(z_{\text{max}}) \simeq 0.3 K_1$ at $z_{\text{max}} \simeq 2.4$. In the weak wash-out regime, when $K_1 \lesssim 3.3$, one has $W_1^{\text{ID}}(z) < 1$ for any value of z . In this case the wash out is negligible and the final asymmetry depends on the initial conditions. In the strong wash-out regime, when $K_1 \gtrsim 3.3$, there is an interval $[z_{\text{on}}, z_{\text{off}}]$ where $W_1^{\text{ID}} \geq 1$. The asymmetry produced at $z \lesssim z_{\text{off}}$ is very efficiently washed out and thus the final asymmetry is essentially what is produced around $z_B \simeq z_{\text{off}}$ by out-of-equilibrium decays of the residual RH neutrinos whose number corresponds approximately to the final value of the efficiency factor.

In Fig. 1 we show the wash-out term from inverse decays for three different values of K_1 . For $K_1 = 100$ we show the interval $[z_{\text{on}}, z_{\text{off}}]$. The maximum value $W_1^{\text{ID}}(z_{\text{max}}) \simeq 33$ is reached at $z_{\text{max}} \simeq 2.4$. For $K_1 \simeq 3.3$ one has $z_{\text{on}} \simeq z_{\text{max}} \simeq z_{\text{off}}$. This can be taken as the threshold value distinguishing between the strong and the weak wash-out regimes. For $K_1 = 10^{-1}$ one has $W_1^{\text{ID}} \ll 1$ for any value of z . Notice however that even in this case the weak wash out can be important for successful leptogenesis if the initial abundance vanishes, since it prevents a full cancellation between two different sign contributions to the final asymmetry [24].

2.2. When are flavor effects important?

We now turn to the question when flavor effects will modify these results. The crucial effect is caused by charged-lepton Yukawa interactions [27] that occur with a rate [28] $\Gamma_{\alpha} \simeq 5 \times 10^{-3} T h_{\alpha}^2$. The largest one is for $\alpha = \tau$ where

$$\frac{\Gamma_{\tau}}{H} \simeq \left(\frac{10^{12} \text{GeV}}{T} \right). \quad (16)$$

Therefore, if $T \gtrsim 10^{12} \text{GeV}$, charged-lepton Yukawa interactions are not effective and all processes in the early Universe are flavor blind, justifying the unflavored treatment. For $T \lesssim 10^{12} \text{GeV}$, the τ Yukawa couplings are strong enough that the scatterings $\tau_L \bar{\tau}_R \rightarrow \Phi^{\dagger}$ are in equilibrium. However, as stressed in the introduction, this condition is not




Figure 1. Comparison between the wash-out term $W_1^{\text{ID}}(z)$ (thick solid lines), defined in Eq. (6) and plotted for the three indicated values of K_1 , and the charged lepton Yukawa interaction term F_τ , defined in Eq. (20) and plotted for the indicated values of M_1 .

necessarily sufficient for important flavor effects to occur because we need to compare the speed of the Yukawa interactions with that of the RH neutrino decays and inverse decays. To this end we study the weak and strong wash-out regimes separately and consider only a two-flavor case because the τ lepton Yukawa coupling causes the main modification.

In the weak wash-out regime, assuming a vanishing initial abundance, the production of RH neutrinos through inverse decays occurs around $T \sim M_1$. At this epoch, inverse decays are by definition slower than the expansion rate. Therefore, the condition $T \lesssim 10^{12}$ GeV is sufficient to conclude that the charged-lepton Yukawa interactions are faster than the inverse decay rate. This translates into the condition $M_1 \lesssim 10^{12}$ GeV because the RH neutrino production occurs at $T \sim M_1$, in agreement with the previous literature [11, 12].

However, this condition does not guarantee that flavor effects indeed have an impact on the final asymmetry, because this impact depends on wash out playing some role. For a vanishing initial abundance this is the case in that wash-out effects prevent a full sign cancellation between the asymmetry produced when $N_{N_1} < N_{N_1}^{\text{eq}}$ and the asymmetry produced later on. On the other hand, for a thermal initial abundance, no such effect arises from the weak wash out and flavor effects do not modify the final asymmetry. In any case, one can say that in the limit $K_1 \rightarrow 0$ flavor cannot have effects on the final asymmetry for any initial abundance. We will come back to this point.

In the strong wash-out regime the situation is very different. The rate of RH neutrino inverse decays at $T \sim M_1$ is larger than the expansion rate. Therefore, we need to compare the charged-lepton Yukawa rate Γ_τ with the RH neutrino inverse decay rate Γ_1^{ID} . For the unflavored treatment to be valid for $z \lesssim z_{\text{fl}} \leq z_B$ then requires

$$M_1 \gtrsim \frac{10^{12} \text{ GeV}}{2 W_1^{\text{ID}}(z_{\text{fl}})}, \quad (17)$$

where z_{fl} is that value of z where the two rates are equal. This condition guarantees that at temperatures $T > T_{\text{fl}} = M_1/z_{\text{fl}}$ flavor effects will not be able to break the coherent propagation of lepton states. The final asymmetry is dominantly produced around $z \sim z_B$. Therefore, the condition for flavor effects to be negligible is

$$M_1 \gtrsim 5 \times 10^{11} \text{ GeV}, \quad (18)$$

similar to the weak wash-out regime. However, the corresponding condition on the temperature

$$T \gtrsim \frac{10^{12} \text{ GeV}}{2 z_B(K_1)} \quad (19)$$

is now less restrictive.

If one starts with a non-vanishing initial abundance, then the final asymmetry is also determined by how efficiently the initial value is washed out; this is described by the integral in Eq. (9). In this case even a value of $z_{\text{fl}} < z_B$ would be important to determine the final asymmetry since wash out in the unflavored regime would be effective for $z \lesssim z_{\text{fl}} < z_B$, while a reduced wash out would apply in the flavored regime at lower temperatures for $z_{\text{fl}} \lesssim z \lesssim z_B$.

We conclude that the condition Eq. (17) obeys the intuitive expectation that there is always a threshold value for K_1 above which the unflavored case is recovered. In this case the temperature below which flavor effects play a role indeed becomes smaller and smaller. The situation is illustrated in Fig. 1 where we compare W_{ID} with

$$F_\tau \equiv \frac{1}{2} \frac{\Gamma_\tau}{H z} \simeq \frac{5 \times 10^{11} \text{ GeV}}{M_1}, \quad (20)$$

the analogous quantity for the charged lepton Yukawa interactions. For any value of M_1 and K_1 , there is a value z_{fl} such that $F_\tau \gtrsim W_{\text{ID}}$ for $z > z_{\text{fl}}$. If $M_1 \lesssim 2 \times 10^{12} \text{ GeV}/K_1$ and $K_1 \gtrsim 3.3$, corresponding to $F_\tau \gtrsim W_{\text{ID}}(z_{\text{max}})$ in the strong wash-out regime, then $z_{\text{fl}} = 0$, meaning that flavor effects are important during the entire thermal history. On the other hand, for a fixed value of M_1 , one has $z_{\text{fl}} \rightarrow \infty$ for $K_1 \rightarrow \infty$, implying that flavor effects tend to disappear for sufficiently large values of K_1 . Notice however that if $M_1 \lesssim 5 \times 10^{11} \text{ GeV}$, then $z_{\text{fl}} \gtrsim z_{\text{off}} \simeq z_B$ for any value of K_1 . This confirms that only for $M_1 \gtrsim 5 \times 10^{11} \text{ GeV}$ flavor effects can be neglected and the unflavored regime is recovered.

3. Maximum flavor effects

We now turn to the opposite extreme case when flavor effects are maximal, the “fully flavored regime.” In other words, the charged-lepton Yukawa interactions are now taken

to be so fast that the lepton flavor content produced in $N \rightarrow \ell + \Phi$ on average fully collapses before the inverse reaction can take place, i.e., the ℓ density matrix in flavor space is to be taken diagonal in the charged-lepton Yukawa basis. In this case each single flavor asymmetry has to be calculated separately because generally the wash out by inverse decays is different for each flavor. Moreover, the single-flavored CP asymmetries now have an additional contribution compared to the total [11, 12]. Finally, the inverse decay involving a lepton in the flavor α does not wash out as much asymmetry as the one produced by one RH neutrino decay. The reduction is quantified by the probability $P_{i\alpha}^0$, averaged over leptons and anti-leptons, that the lepton ℓ_i produced in the decay of N_i collapses into the flavor eigenstate ℓ_α . The relevant Boltzmann equations become

$$\begin{aligned} \frac{dN_{N_1}}{dz} &= -D_1 (N_{N_1} - N_{N_1}^{\text{eq}}) \\ \frac{dN_{\Delta_\alpha}}{dz} &= \varepsilon_{1\alpha} D_1 (N_{N_1} - N_{N_1}^{\text{eq}}) - P_{1\alpha}^0 W_1^{\text{ID}} N_{\Delta_\alpha}. \end{aligned} \quad (21)$$

Since we are dealing with the two-flavor case, here $\alpha = \tau$ or μ where the latter stands for a suitable superposition of the μ and e flavor. Notice also that we defined $\Delta_\alpha \equiv B/2 - L_\alpha$ and therefore the total asymmetry is given by $N_{B-L} = N_{\Delta_\mu} + N_{\Delta_\tau}$.

As in the unflavored case, we next identify the condition for the fully flavored approximation to hold. The final asymmetry in the flavor α is dominantly produced at $z \simeq z_{B\alpha} \equiv z_B(K_{1\alpha})$, where $K_{1\alpha} \equiv P_{1\alpha}^0 K_1$. Therefore, one must require that $\Gamma_\alpha \gtrsim \Gamma_1^{\text{ID}}$ holds already at $z \sim z_{B\alpha}$ lest the wash-out reduction takes place too late. We stress that flavor effects modify the final asymmetry only if the flavor projection takes place before the wash out by inverse decays freezes out. Otherwise the wash-out epoch is over and the unflavored behavior is recovered. It is easy to verify that if the projectors are set to unity and the equations are summed over flavors, the kinetic equation for N_{B-L} holding in the unflavored regime is recovered [cf. Eq. (3)]. Therefore, we require

$$M_1 \lesssim \frac{10^{12} \text{ GeV}}{2 W_1^{\text{ID}}(z_{B\alpha})} \quad (22)$$

as an approximate condition for the fully flavored behavior.

In Fig. 2 we summarize the different possible cases in the plane of parameters K_1 and M_1 . For $M_1 \gtrsim 5 \times 10^{11} \text{ GeV}$, above the dashed line, flavor effects are not important independently of K_1 . The condition Eq. (22), in the most restrictive case when $z_{B\alpha} = z_{\text{max}}$ and $W_1^{\text{ID}} \simeq 0.3 K_1$, is satisfied below the inclined dotted line. This case typically occurs in a one flavor dominated scenario, as we explain below. The vertical dot-dashed line is the border that separates the weak from the strong wash-out regime in the unflavored case. In the flavored case the condition $K_1 \lesssim 3.3$ still implies the weak wash-out regime because flavor effects can only reduce the wash out. However, the condition for the strong wash-out regime can be more restrictive than $K_1 \gtrsim 3.3$, as discussed in [14]. For $K_1 \lesssim 3.3$, flavor effects modify the final asymmetry only marginally and more specifically only if the initial abundance vanishes, as indicated in Fig. 2. On the other hand, for $K_1 \gtrsim 3.3$ and below the diagonal line, flavor modifications of the final asymmetry can be large, especially in the one-flavor dominated scenario.




Figure 2. Relevance of flavor effects in schematic regions of parameters K_1 and M_1 . The region above the horizontal dashed line corresponds to the condition (18) for $z_{B\alpha} = z_{\max}$. The vertical dot-dashed line is the border between the weak and the strong wash-out regime. The region below the inclined dotted line corresponds to the condition (22) for $z_{B\alpha} = z_{\max}$.

There is a region in parameter space where neither condition Eq. (17) nor (22) holds. This intermediate regime can become very large in the case of a one-flavor dominated scenario, where an order of magnitude enhancement of the final asymmetry is possible. In this case one of the two projectors is very small compared to the other and so the wash out is very asymmetric in the two flavors. On the other hand, if the two-flavored CP asymmetries are comparable, then the final asymmetry is dominantly produced into one flavor and deviations from the unflavored regime can become very large. This scenario is realized, in particular, when the absolute neutrino mass scale increases, relaxing the traditional neutrino mass bound.

However, the condition Eq. (22) strongly restricts the applicability of the one-flavor dominated scenario. Even though the reduction of the wash out is driven by $K_{1\alpha} \ll K_1$, implying $z_{B\alpha} \ll z_B$, the possibility for flavor effects to be relevant relies on the dominance of the charged-lepton Yukawa interaction rate compared to the RH neutrino inverse decay rate that however is still driven by K_1 . Therefore, increasing K_1 , one can enhance the asymmetry in the one-flavor dominated scenario compared to the unflavored case, if smaller and smaller values of the projector $P_{1\alpha}^0$ are possible. On the other hand, the inverse decay rate increases so that the fully flavored behavior may no longer apply. In particular notice that the maximum enhancement of the asymmetry is obtained when $K_1 \gg K_{1\alpha} \simeq 1$, when $z_{B\alpha} \simeq z_{\max}$ and the condition (22) is maximally restrictive.

4. Neutrino mass bound

One possible consequence of flavor effects is to relax the traditional upper bound on the neutrino mass that is implied by successful leptogenesis. In order to explore the impact of our modified criteria we first recall the origin of this bound in the unflavored case. Maximizing the final value of the asymmetry over all see-saw parameters except M_1 and m_1 yields [24]

$$\frac{\eta_B^{\max}(M_1, m_1)}{\eta_B^{\text{CMB}}} \simeq 3.8 \left(\frac{m_\star}{m_1}\right)^{1.2} \left(\frac{M_1}{10^{10} \text{ GeV}}\right) \frac{m_{\text{atm}}}{m_1 + m_3} \exp \left[-\frac{w}{z_B} \frac{M_1}{10^{10} \text{ GeV}} \left(\frac{\bar{m}}{\text{eV}}\right)^2 \right] \geq 1, \quad (23)$$

where we have approximated $\kappa(K_1) \simeq 0.5 K_1^{-1.2}$ [29] and we have neglected the dependence of z_B on K_1 in the derivative. This constraint translates into $m_1 < m_1^{\max}(M_1)$ shown by the curved solid line in the upper part of Fig. 3 where the unflavored behavior obtains. This curve sports an absolute maximum, $m_1 \lesssim 0.12 \text{ eV}$, for $M_1 \simeq 10^{13} \text{ GeV}$.

The possibility that flavor effects could relax this bound is based on the observation that the flavored CP asymmetries, for $m_1 \gtrsim m_{\text{atm}}$, are proportional to m_1 . On the other hand, the total asymmetry is suppressed like m_1^{-1} , contributing to the upper bound in the unflavored regime. However, if m_1 increases, then K_1 has to increase also so that it is not guaranteed that the fully flavored treatment remains justified.

Quantitatively, the value of K_1 is bounded from below by [30]

$$K_1 \geq \frac{m_1}{m_\star}, \quad (24)$$

where

$$m_\star = \frac{16 \pi^{5/2} \sqrt{g_\star}}{3 \sqrt{5}} \frac{v^2}{M_{\text{Pl}}} \simeq 1.08 \times 10^{-3} \text{ eV}. \quad (25)$$

For $K_1 \geq m_1/m_\star \gg 1$, the final asymmetry is maximized when a one-flavor dominated scenario is realized. In this case the final asymmetry is approximately $N_{B-L}^{\text{f}} \simeq \varepsilon_{1\alpha} \kappa_{1\alpha}^{\text{f}}$. The bound on the single flavored CP asymmetry is [13]

$$|\varepsilon_{1\alpha}| < \bar{\varepsilon}(M_1) \sqrt{P_{1\alpha}^0} \frac{m_3}{m_{\text{atm}}}, \quad (26)$$

where

$$\bar{\varepsilon}(M_1) \equiv \frac{3}{16\pi} \frac{M_1 m_{\text{atm}}}{v^2} \simeq 10^{-6} \left(\frac{M_1}{10^{10} \text{ GeV}}\right) \left(\frac{m_{\text{atm}}}{0.05 \text{ eV}}\right). \quad (27)$$

It is then possible to find the value of $P_{1\alpha}^0$ that maximizes the asymmetry as a function of K_1 and the corresponding value of $z_{B\alpha}$. Imposing $\eta_B^{\max} \geq \eta_B^{\text{CMB}}$ implies a lower bound on M_1 as a function of K_1 .

This limit can be translated into a lower bound on M_1 as a function of m_1 by replacing K_1 with its minimum value $m_1/m_\star$. In this way the wash out is always minimized and the final efficiency factor and the final asymmetry are maximized. Notice that the single-flavored CP asymmetries, like the total, vanish for $K_1 = m_1/m_\star$.

Therefore, this lower bound cannot be saturated. Notice moreover that for $m_1 \lesssim m_*$ the one-flavor dominated scenario does not necessarily holds because it is possible that $K_1 \lesssim 1$. Actually for $K_1 \rightarrow 0$, flavor effects disappear and one has to recover the usual asymptotic value of the lower bound obtained in the unflavored case for thermal initial abundance, $M_1 \gtrsim 4 \times 10^8 \text{ GeV}$. For intermediate values of K_1 one can use a simple interpolation. The final result is shown in the bottom part of Fig. 3 as a thick solid line.

In Fig. 3 we also show the condition Eq. (22), calculated for the same value of $z_{B\alpha}$ that maximizes the final asymmetry, but replacing K_1 with its minimum value m_1/m_* (thick dashed line). Since W_1^{ID} increases with K_1 , this produces a necessary, but not sufficient, condition in the m_1 - M_1 plane for the fully flavored behavior. This condition




Figure 3. Relevance of flavor effects similar to Fig. 2, now mapped to schematic regions of parameters m_1 and M_1 . The region above the horizontal dashed line corresponds to the condition Eq. (18) for the applicability of the unflavored regime. The region below the inclined thick dashed line corresponds to the condition Eq. (22) calculated for that value of $z_{B\alpha}$ that maximizes the final asymmetry in the one-flavor dominated scenario and for $K_1 = m_1/m_*$. In this same case, the lower inclined dot-dashed lines includes also the effect of scatterings in the condition Eq. (22) while the upper inclined and the horizontal dot-dashed lines include the effect of oscillations. The area between the two inclined dot-dashed lines gives an estimation of the uncertainty on the condition for the fully flavored regime to hold. The area between the horizontal thin dot-dashed line and the horizontal thick dashed line gives an estimation of the uncertainty on the condition for the unflavored regime to hold. The two thick solid lines borders the region where successful leptogenesis is possible: on the left in the unflavored regime and above in the fully flavored regime. The thin solid line is a more restrictive border obtained for a specific choice of the see-saw orthogonal matrix and the thin dashed line is the corresponding condition Eq. (22). In this case one has $K_1 > m_1/m_*$.

matches the validity of the unflavored regime at $m_1 \simeq 3 \times 10^{-3}$ eV and the lower bound on M_1 at $m_1 \simeq 2$ eV. This means that for $m_1 \gtrsim 2$ eV the fully flavored behavior does not obtain. Notice also that this upper limit is quite conservative because the lower bound on M_1 has been obtained neglecting that, for $K_1 = m_1/m_*$, the flavored CP asymmetry vanishes and thus the bound cannot be saturated. Moreover we have assumed that $P_{1\alpha}^0$ can always assume the value that maximizes the asymmetry.

In Fig. 3 we also show (thin solid line) a lower bound $M_1(m_1)$ in a specific scenario [14], corresponding to a particular choice of the orthogonal see-saw matrix [31] $\Omega = R_{13}$, that represents a complex rotation in the 13-plane, where Ω_{13}^2 is taken purely imaginary. In this case the value of $z_{B\alpha}$ is not necessarily the same that maximizes the asymmetry in the one-flavor dominated scenario and $K_1 > m_1/m_*$. Therefore, a specific calculation is necessary in order to work out correctly the condition Eq. (22). The result is shown in Fig. 3 with a thin dashed line.

In this case the upper limit on m_1 for the applicability of the fully flavored regime is much smaller, $m_1 \simeq 0.1$ eV. Allowing for a non-vanishing real part of Ω_{13}^2 , slightly larger values are possible. It should be however kept in mind that these values are indicative since they rely on a condition for the fully flavored regime that comes from a simple rate comparison.

5. Limitations of a simple rate comparison

We have exploited a somewhat qualitative rate comparison for the determination of the region where the fully flavored regime obtains. While we believe that our approach nicely illustrates the modifications that derive from our more restrictive criterion for the significance of flavor effects, there are also important shortcomings. First, we have simply compared the inverse decay rate with the charged-lepton Yukawa interaction rate, ignoring flavor oscillations caused by the flavor-dependent lepton dispersion relation in the medium. If the oscillations are much faster than the inverse decay rate, they also contribute effectively, together with inelastic scatterings, to project the lepton state on the flavor basis. Therefore, including oscillations will tend to enlarge the region where the fully flavored behavior obtains (the inclined upper dot-dashed line in Fig. 3) and to reduce that one where the unflavored behavior obtains (the horizontal upper dot-dashed line in Fig. 3). In our case, the oscillation frequency is comparable to Γ_α and so the two estimations are not too far off.

Moreover, we have also neglected $\Delta L = 1$ scatterings. They also contribute, like inverse decays, both to generate the asymmetry and to wash out and therefore, together with inverse decays, contribute to preserving the flavor direction of the leptons. At the relevant $z \sim z_{B\alpha} \sim 2$, the $\Delta L = 1$ scattering rate is actually larger than the inverse decay rate and thus tends to reduce the region where the fully flavored behavior obtains (the lower inclined dot-dashed line in Fig. 3). Therefore, the effects of oscillations and of $\Delta L = 1$ scatterings may partially cancel each other. In Fig. 3 the region between the two inclined dot-dashed lines gives therefore an indication of

the theoretical uncertainty on the determination of the region where the fully flavored regime holds. It can be seen that current calculations cannot establish whether the upper bound holding in the unflavored regime is nullified, just simply relaxed or still holding, when flavor effects are included.

Only a full quantum-kinetic treatment can give a final verdict on the effectiveness of flavor effects in leptogenesis and its impact on the neutrino mass limit. While we have verified that our rate criteria are borne out by the quantum kinetic equations stated in Ref. [12], these equations are not necessarily complete in that the terms describing the generation and wash out of the asymmetry have been added by hand. Moreover, the “damping rate” caused by the flavor-sensitive Yukawa interactions ultimately derives from a collision term in the kinetic equation [18]. Extending the pioneering treatment of Ref. [12] to allow for a complete understanding of flavor effects remains a challenging task.

6. Conclusions

Flavor effects can play a very important role in leptogenesis. However, we have shown that the condition for the fully flavored behavior is more restrictive because one needs to compare the speed of the charged-lepton Yukawa interactions with that of $N \leftrightarrow \ell + \Phi$, not with the cosmic expansion rate. This distinction makes a significant difference particularly in the strong wash-out regime where for some of the leptogenesis epoch the rate of $N \leftrightarrow \ell + \Phi$ is faster than the cosmic expansion.

We are especially interested in the question if the traditional neutrino mass bound of the unflavored treatment [8] can be circumvented by flavor effects. We have found that the see-saw parameters that correspond to $m_1 \gtrsim 0.1$ eV do not fall into the strictly unflavored or the fully flavored regimes. The intermediate regime requires a detailed quantum kinetic treatment, so at present it is not possible to decide if the upper bound neutrino mass bound can be circumvented (relaxed or nullified) by flavor effects.

Acknowledgments

It is a pleasure to thank W. Buchmüller, S. Davidson, M. Losada, M. Plümacher and E. Roulet for comments and discussions. We also wish to thank A. Riotto for sending us the draft of a forthcoming manuscript with A. De Simone on the same subject. This work was supported, in part, by the Deutsche Forschungsgemeinschaft (DFG) under grant No. SFB-375 and by the European Union under the ILIAS project, contract No. RII3-CT-2004-506222, and under the Marie Curie project “Leptogenesis, Seesaw and GUTs,” contract No. MEIF-CT-2006-022950.

References

- [1] P. Minkowski, “ $\mu \rightarrow e\gamma$ at a rate of one out of 10^9 muon decays?,” *Phys. Lett. B* **67** (1977) 421.
- [2] T. Yanagida, “Horizontal gauge symmetry and masses of neutrinos,” in *Proc. Workshop on the Baryon Number of the Universe and Unified Theories* (13–14 Feb 1979, Tsukuba, Japan), p. 95.
- [3] M. Gell-Mann, P. Ramond and R. Slansky, “Complex spinors and unified theories,” in *Supergravity*, eds. P. van Nieuwenhuizen and D. Freedman (North Holland, Amsterdam, 1979) p. 315.
- [4] S. L. Glashow, “The future of elementary particle physics,” in *1979 Cargese Summer Institute on Quarks and Leptons*, eds. M. Levy, J.-L. Basdevant, D. Speiser, J. Weyers, R. Gastmans and M. Jacobs, (Plenum Press, New York, 1980), NATO Adv. Study Inst. Ser. B Phys. **59** (1979) 687.
- [5] R. Barbieri, D. V. Nanopoulos, G. Morchio and F. Strocchi, “Neutrino masses in grand unified theories,” *Phys. Lett. B* **90** (1980) 91.
- [6] R. N. Mohapatra and G. Senjanović, “Neutrino mass and spontaneous parity nonconservation,” *Phys. Rev. Lett.* **44** (1980) 912.
- [7] M. Fukugita and T. Yanagida, “Baryogenesis without grand unification,” *Phys. Lett. B* **174** (1986) 45.
- [8] W. Buchmüller, P. Di Bari and M. Plümacher, “The neutrino mass window for baryogenesis,” *Nucl. Phys. B* **665** (2003) 445 [hep-ph/0302092].
- [9] R. Barbieri, P. Creminelli, A. Strumia and N. Tetradis, *Nucl. Phys. B* **575** (2000) 61 [arXiv:hep-ph/9911315].
- [10] T. Endoh, T. Morozumi and Z. h. Xiong, “Primordial lepton family asymmetries in seesaw model,” *Prog. Theor. Phys.* **111** (2004) 123 [hep-ph/0308276].
- [11] E. Nardi, Y. Nir, E. Roulet and J. Racker, “The importance of flavor in leptogenesis,” *JHEP* **0601** (2006) 164 [hep-ph/0601084].
- [12] A. Abada, S. Davidson, F. X. Josse-Michaux, M. Losada and A. Riotto, “Flavour issues in leptogenesis,” *JCAP* **0604** (2006) 004 [hep-ph/0601083].
- [13] A. Abada, S. Davidson, A. Ibarra, F. X. Josse-Michaux, M. Losada and A. Riotto, “Flavour matters in leptogenesis,” *JHEP* **0609** (2006) 010 [hep-ph/0605281].
- [14] S. Blanchet and P. Di Bari, “Flavor effects on leptogenesis predictions,” hep-ph/0607330.
- [15] S. Pascoli, S. T. Petcov and A. Riotto, “Connecting low energy leptonic CP-violation to leptogenesis,” hep-ph/0609125.
- [16] G. C. Branco, R. González Felipe and F. R. Joaquim, “A new bridge between leptonic CP violation and leptogenesis,” hep-ph/0609297.
- [17] L. Stodolsky, “On the treatment of neutrino oscillations in a thermal environment,” *Phys. Rev. D* **36** (1987) 2273.
- [18] G. Raffelt, G. Sigl and L. Stodolsky, “Non-Abelian Boltzmann equation for mixing and decoherence,” *Phys. Rev. Lett.* **70** (1993) 2363 [hep-ph/9209276].
- [19] P. Di Bari, “Seesaw geometry and leptogenesis,” *Nucl. Phys. B* **727** (2005) 318 [hep-ph/0502082].
- [20] O. Vives, “Flavoured leptogenesis: A successful thermal leptogenesis with $N(1)$ mass below $10^8 - \text{GeV}$,” *Phys. Rev. D* **73** (2006) 073006 [hep-ph/0512160].
- [21] G. F. Giudice, A. Notari, M. Raidal, A. Riotto and A. Strumia, “Towards a complete theory of thermal leptogenesis in the SM and MSSM,” *Nucl. Phys. B* **685** (2004) 89 [hep-ph/0310123].
- [22] W. Buchmüller and M. Plümacher, “Spectator processes and baryogenesis,” *Phys. Lett. B* **511** (2001) 74 [hep-ph/0104189].
- [23] E. Nardi, Y. Nir, J. Racker and E. Roulet, “On Higgs and sphaleron effects during the leptogenesis era,” *JHEP* **0601** (2006) 068 [hep-ph/0512052].
- [24] W. Buchmüller, P. Di Bari and M. Plumacher, “Leptogenesis for pedestrians,” *Annals Phys.* **315** (2005) 305 [hep-ph/0401240].
- [25] S. Blanchet and P. Di Bari, “Leptogenesis beyond the limit of hierarchical heavy neutrino masses,” *JCAP* **0606** (2006) 023 [hep-ph/0603107].

- [26] D. N. Spergel *et al.*, “Wilkinson Microwave Anisotropy Probe (WMAP) three year results: Implications for cosmology,” astro-ph/0603449.
- [27] R. Barbieri, P. Creminelli, A. Strumia and N. Tetradis, “Baryogenesis through leptogenesis,” Nucl. Phys. B **575** (2000) 61 [hep-ph/9911315].
- [28] B. A. Campbell, S. Davidson, J. R. Ellis and K. A. Olive, “On the baryon, lepton flavor and right-handed electron asymmetries of the universe,” Phys. Lett. B **297** (1992) 118 [hep-ph/9302221].
- [29] P. Di Bari, “Leptogenesis, neutrino mixing data and the absolute neutrino mass scale,” arXiv:hep-ph/0406115.
- [30] M. Fujii, K. Hamaguchi and T. Yanagida, “Leptogenesis with almost degenerate Majorana neutrinos,” Phys. Rev. D **65** (2002) 115012 [arXiv:hep-ph/0202210].
- [31] J. A. Casas and A. Ibarra, “Oscillating neutrinos and $\mu \rightarrow e, \gamma$,” Nucl. Phys. B **618** (2001) 171 [hep-ph/0103065].